PROJECT NUMBER: NASA-WASHINGTON PROJECT- NAGW 1998

TITLE: NASA SUPPORTING STUDIES FOR MICROGRAVITY RESEARCH ON EYE MOVEMENTS- FINAL TECHNICAL REPORT

PRINCIPAL INVESTIGATOR: BERNARD COHEN, M.D.

INSTITUTION: MOUNT SINAI SCHOOL OF MEDICINE

DATES OF PROJECT: 1/1/90 - 12/31/90

ADDRESS OF P.I.:

Dr. Bernard Cohen
Department of Neurology, Box 1135
Mount Sinai School of Medicine
1 East 100th Street
New York, New York 10029
Telephone: 212-241-7068

FAX: 212-831-1610

FINAL REPORT:

The purpose of the work on this project was to provide support for ground-based studies on the effects of gravity on eye movements. In this research we investigated the efffects of microgravity on the optokinetic eye movements of humans. OKN was induced by having subjects watch 3.3° stripes moving at 35°/s for 45 s in a binocular, head-fixed apparatus. The field (hor., 88°; vert., 72°), was rotated about axes that were upright or tilted 45° or 90°. The head was upright or tilted 45° on the body. Head-horizontal (yaw axis) and head-vertical (pitch axis) components of OKN were recorded with electro-oculography (EOG). Slow phase velocity vectors were determined relative to gravity. With the head upright, the axis of eye rotation during yaw axis OKN was coincident with the stimulus axis and the spatial vertical. With the head tilted 45° on the body, a persistent vertical component of eye velocity developed during yaw axis stimulation, and there was an average shift of the axis of eye rotation toward the spatial vertical of ≈ 18° in six subjects. During oblique optokinetic stimulation with the head upright, the axis of eye rotation shifted 12° toward the spatial vertical. When the head was tilted, the axis of eye rotation rotated to the other side of the spatial vertical by 5.4° during the same oblique stimulation. This counter-rotation of the axis of eye rotation is similar to the "Müller (E) effect", in which the perception of the upright counter-rotates to the opposite side of the spatial vertical when subjects are tilted in darkness.

The data were simulated by a model of OKN. Despite the short OKAN time constants, strong horizontal to vertical cross-coupling was produced if the horizontal and vertical time constants were in proper ratio, and there was no suppression of nystagmus orthogonal to the stimulus direction. This shows that the spatial orientation of OKN can be due to a restructuring of the system matrix of velocity storage as a function of gravity. We conclude that although human OKAN is weak, velocity storage orients the slow phase velocity of OKN towards the spatial vertical.

A paper describing these results has been submitted to Experimental Brain Research.

FINAL REPORT:

The purpose of the work on this project was to provide support for ground-based studies on the effects of gravity on eye movements. In this research we investigated the efffects of microgravity on the optokinetic eye movements of humans. OKN was induced by having subjects watch 3.3° stripes moving at 35% for 45 s in a binocular, head-fixed apparatus. The field (hor., 88°; vert., 72°), was rotated about axes that were upright or tilted 45° or 90°. The head was upright or tilted 45° on the body. Head-horizontal (yaw axis) and head-vertical (pitch axis) components of OKN were recorded with electro-oculography (EOG). Slow phase velocity vectors were determined relative to gravity. With the head upright, the axis of eye rotation during yaw axis OKN was coincident with the stimulus axis and the spatial vertical. With the head tilted 45° on the body, a persistent vertical component of eye velocity developed during yaw axis stimulation, and there was an average shift of the axis of eye rotation toward the spatial vertical of ≈ 18° in six subjects. During oblique optokinetic stimulation with the head upright, the axis of eye rotation shifted 12° toward the spatial vertical. When the head was tilted, the axis of eye rotation rotated to the other side of the spatial vertical by 5.4° during the same oblique stimulation. This counter-rotation of the axis of eye rotation is similar to the "Müller (E) effect", in which the perception of the upright counter-rotates to the opposite side of the spatial vertical when subjects are tilted in darkness.

The data were simulated by a model of OKN. Despite the short OKAN time constants, strong horizontal to vertical cross-coupling was produced if the horizontal and vertical time constants were in proper ratio, and there was no suppression of nystagmus orthogonal to the stimulus direction. This shows that the spatial orientation of OKN can be due to a restructuring of the system matrix of velocity storage as a function of gravity. We conclude that although human OKAN is weak, velocity storage orients the slow phase velocity of OKN towards the spatial vertical.

A paper describing these results has been submitted to Experimental Brain Research.

June 23, 1993

Dr. Janis Stoklosa
Mail Code ULM, Office 2441
Life and Microgravity Sciences
and Applications
NASA Headquarters
Washington DC 20546-0001

Dear Dr. Stoklosa:

Please find enclosed the final technical report for our project NASA-Washinton NAGW 1998, NASA SUPPORTING STUDIES FOR MICROGRAVITY RESEARCH ON EYE MOVEMENTS.

Sincerely,

Bernard Cohen, M.D.

ONIE No. 11990-16081 Rapiretion Date: 12/31/

Ş.€ FOR GOV VOUCHER NO. PAGE A 2. PUBLIC VOUCHER NO. 18, DATE REC'D, MOVYR DATE 4. REPORT NO. Pico Pico ACCEPTED BY AUTHORIZED WAY HIMSENT ₽ 00. NO. NAGW 1988 7. NAME AND ADDRESS OF CONTRACTOR 3. CONTRACT MUNBER UNT ACQUISITION SIGNATURE AND TITLE MFR. SERIAL ć C ACQUISITION—GOV, TITLED

ACQUISITION—CONTR, TITLED

ANNUAL INVENTORY

FINAL INVENTORY MODEL OR report of **can**talized monexpendable equapment 19, AUTHENTICATION BY CONTRACTOR'S SUPERVISORY ACCOUNTING OFFICIAL NOTE: Complete this form in accordance with instructions on means eige of copy Ma. 6 4. TYPE OF REPORT HEALTH, AND HUMAN SERVICES DATE DEPARTMENT OF ž 15 0 E a name of Person responsible for this report BURNARD CONON DESCRIPTION & HEN 212 241 7068 NAME (TYPED) HHG 565 (7/70) SIGNATURE ITEM NO.